

Scotland's Rural College

## First isolation of *Arcanobacterium pinnipediorum* from a grey seal pup (*Halichoerus grypus*) in the UK

Alssahen, Mazen; Foster, Geoffrey; Hassan, Abdulwahed Ahmed; Rau, Jörg; Lämmle, Christoph; Prenger-Berninghoff, Ellen; Eisenberg, Tobias; Robinson, Mathew; Abdulmawjood, Amir

*Published in:*  
*Folia Microbiologica*

DOI:  
[10.1007/s12223-021-00932-7](https://doi.org/10.1007/s12223-021-00932-7)

Print publication: 01/04/2022

*Document Version*  
Publisher's PDF, also known as Version of record

[Link to publication](#)

### *Citation for published version (APA):*

Alssahen, M., Foster, G., Hassan, A. A., Rau, J., Lämmle, C., Prenger-Berninghoff, E., Eisenberg, T., Robinson, M., & Abdulmawjood, A. (2022). First isolation of *Arcanobacterium pinnipediorum* from a grey seal pup (*Halichoerus grypus*) in the UK. *Folia Microbiologica*, 67(2), 291-297. <https://doi.org/10.1007/s12223-021-00932-7>

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

### Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



# First isolation of *Arcanobacterium pinnipediorum* from a grey seal pup (*Halichoerus grypus*) in the UK

Mazen Alssahen<sup>1</sup> · Geoffrey Foster<sup>2</sup> · Abdulwahed Ahmed Hassan<sup>3</sup> · Jörg Rau<sup>4</sup> · Christoph Lämmle<sup>1</sup> · Ellen Prenger-Berninghoff<sup>1</sup> · Tobias Eisenberg<sup>1,5</sup> · Mathew Robinson<sup>6</sup> · Amir Abdulmawjood<sup>7</sup>

Received: 30 March 2021 / Accepted: 8 November 2021  
© The Author(s) 2021

## Abstract

In the present study, a single *Arcanobacterium* (*A.*) *pinnipediorum* strain isolated from discharge of a jaw swelling of a grey seal pup (*Halichoerus grypus*) in England, UK, was identified. This strain was further characterized by phenotypical investigations, by matrix-assisted laser desorption/ionization time-of-flight mass spectrometry (MALDI-TOF MS), by Fourier transform infrared spectroscopy (FT-IR), and genotypically by sequencing the 16S rRNA gene and the genes *gap* encoding glyceraldehyde 3-phosphate dehydrogenase, *tuf* encoding elongation factor tu, and *rpoB* encoding the  $\beta$  subunit of bacterial RNA polymerase. The present study gives a first detailed characterization of the species *A. pinnipediorum* from a grey seal in the UK. However, the route of infection of the grey seal with the bacterial pathogen remains unclear.

## Introduction

The genus *Arcanobacterium* (*A.*) was described by Collins et al. (1982) as a group of facultative anaerobic, asporogenous Gram-positive rods. According to Yassin et al. (2011) genus *Arcanobacterium* consists of four species, namely, *A. haemolyticum*, *A. phocae*, *A. pluranimalium*, and *A. hippocoleae*. In the following years, *A. canis* (Hijazin et al. 2012b), *A.*

*phocisimile* (Hijazin et al. 2013), *A. pinnipediorum* (Sammra et al. 2015), *A. wilhelmae* (Sammra et al. 2017), *A. urinimassiliense* (Diop et al. 2017), *A. ihumii* (Fall et al. 2019), and *A. bovis* (Sammra et al. 2020) were described as novel species of this genus. The type strain *A. pinnipediorum* DSM 2710<sup>T</sup> was originally isolated in 2004 during a monitoring program from an anal swab of a female harbor seal and characterized phenotypically, by cell wall analysis and genotypically (Sammra et al. 2015, 2018). To the best of our knowledge, no other *A. pinnipediorum* strains have been described elsewhere. The present study was designed to characterize a single *A. pinnipediorum* strain, which represents the first isolation of this bacterial species in the UK and the first from a grey seal (*Halichoerus grypus*).

## Materials and methods

The bacterial strains investigated in the present study included *A. pinnipediorum* 014418, type strain *A. pinnipediorum* DSM 28752<sup>T</sup>, originally isolated in the German North Sea, the type strains *A. phocae* DSM 10002<sup>T</sup> and *A. phocisimile* DSM 26142<sup>T</sup>, and other type strains of genus *Arcanobacterium* (Hijazin et al. 2012b, 2013; Sammra et al. 2015). *A. pinnipediorum* 014418 was isolated in mixed culture from a swollen mandible discharge of an 8-month-old male grey seal pup in a rehabilitation centre in England. Upon submission, the seal had a high temperature, infected

✉ Mazen Alssahen  
Mazen.Alssahen@vetmed.uni-giessen.de

<sup>1</sup> Institut Für Hygiene Und Infektionskrankheiten Der Tiere, Justus-Liebig-Universität Giessen, Frankfurter Straße 85-91, Giessen 35392, Germany

<sup>2</sup> SRUC Veterinary Services, 10 Inverness Campus, Inverness, An Lochran IV2 5NA, UK

<sup>3</sup> Department of Veterinary Public Health (DVPH), College of Veterinary Medicine, Mosul University, Mosul, Iraq

<sup>4</sup> Chemisches Und Veterinäruntersuchungsamt Stuttgart (CVUAS), Schaflandstraße 3/2, Fellbach 70736, Germany

<sup>5</sup> Landesbetrieb Hessisches Landeslabor, Giessen, Schubertstraße 60, Giessen 35392, Germany

<sup>6</sup> Axiom Veterinary Laboratories, Manor House, Brunel Road, Newton Abbot TQ12 4PB, UK

<sup>7</sup> Institute of Food Quality and Food Safety, Research Center for Emerging Infections and Zoonoses (RIZ), University of Veterinary Medicine Hannover, Bischofsholer Damm 15, Hannover 30173, Germany

teeth, and gums with discharging pus around the teeth. The animal was treated with amoxicillin/clavulanic acid and clindamycin for 17 days and the discharge resolved. Six weeks later a swab was collected from the bilateral anterior mandible that was swelling and discharging leading to the isolation of *A. pinnipediorum* 014418. The animal was treated again with amoxicillin/clavulanic acid and clindamycin for 25 days. In addition, chin became a hard swelling but there was no discharge. The animal was released 10 days later.

The bacterial culturing of *A. pinnipediorum* 014418 was carried out on 5% sheep blood agar for 48 h at 37 °C under microaerobic conditions. The strain was investigated phenotypically by determination of hemolysis and by VITEK2-compact system (bioMérieux, Nürtingen, Germany) according to the instructions of the manufacturer. The *A. pinnipediorum* strain was additionally analyzed by matrix-assisted laser desorption ionization-time of flight mass spectrometry (MALDI TOF MS, Bruker Biotyper database 8.468, Bruker Daltonik, Bremen, Germany) (Sammra et al. 2018; Wickhorst et al. 2019) and by Fourier transform infrared spectroscopy (FT-IR, Bruker Tensor with HTS-XT, Bruker Optik, Ettlingen, Germany), (Nagib et al. 2014; Sammra et al. 2018).

The presence of *A. phocae* phocaelysin encoding gene *phl* was determined with a previously described loop-mediated isothermal amplification (LAMP) assay. This was performed using a real-time fluorometer (Genie II®, OptiGene, Horsham, UK) (Abdulmawjood et al. 2016; Alssahen et al. 2018).

*A. pinnipediorum* 014418 was additionally investigated by amplification and sequencing of 16S rRNA gene (Hassan et al. 2009; Sammra et al. 2014b, 2015). The DNA template was extracted from freshly cultivated bacterial colonies using the DNeasy Blood and Tissue Kit in accordance

with the manufacturer's instructions (Qiagen, Hilden, Germany). The nucleotide sequence of the bacterial 16S rRNA gene was amplified with forward primer 16S rRNA UNI-L (5'-AGA GTT TGA TCA TGG CTC AG-3') and reverse primer 16S rRNA UNI-R (5'-GTG TGA CGG GCG GTG TGT AC-3'). The amplicon was used for sequencing with primer 16S rRNA-533F (5'-GTG CCA GCM GCC GCG GTA A-3') and 16S rRNA-907-R (5'-CCG TCA ATT CMT TTG AGT TT-3') by Eurofins Genomics GmbH (Ebersberg, Germany). The sequences were analyzed using FinchTV (version, 1.4.0), alignment, and phylogenetic analysis by the Clustal W method using DNASTAR Lasergene version 8.0.2 (DNASTAR Inc., Madison, USA). Furthermore, *A. pinnipediorum* 014418 was characterized by sequencing the genes *gap* encoding glyceraldehyde 3-phosphate dehydrogenase, *tuf* encoding elongation factor tu (Wickhorst et al. 2017, 2019; Sammra et al. 2018), and *rpoB* encoding the  $\beta$  subunit of bacterial RNA polymerase (Ülbeği-Mohyla et al. 2010; Sammra et al. 2013, 2018). The oligonucleotide primer sequences and PCR conditions of the target genes used in the present study are summarized in Table 1.

## Results and discussion

*A. pinnipediorum* 014418 investigated in the present study displayed a relatively small zone of complete hemolysis on 5% sheep blood agar plates and could be classified biochemically using VITEK2-compact system. The biochemical properties of *A. pinnipediorum* 014418 mainly corresponded to properties of *A. pinnipediorum* DSM 28752<sup>T</sup>. However, differences were observed in a positive D-galactose reaction and negative beta-D-fucosidase reaction of *A. pinnipediorum* 014418 (Table 2). Strain *A. pinnipediorum*

**Table 1** Oligonucleotide primer sequences and PCR conditions of the target genes used in the present study

Oligonucleotide primer	Sequence	Expected size of PCR product (bp)	Program*
16S rRNA UNI-L	5'-AGA GTT TGA TCA TGG CTC AG-3'	1403	1
16S rRNA UNI-R (amplification primer)	5'-GTG TGA CGG GCG GTG TGT AC-3'		
16S rRNA-533-F	5'-GTG CCA GCM GCC GCG GTA A-3'	—	—
16S rRNA-907-R (sequencing primer)	5'-CCG TCA ATT CMT TTG AGT TT-3'		
Gap-F	5'-TCG AAG TTG TTG CAG TTA ACG A-3'	830	2
Gap-R	5'-CCA TTC GTT GTC GTA CCA AG-3'		
Tuf-F	5'-GGA CGG TAG TTG GAG AAG AAT GG-3'	796	3
Tuf-R	5'-CCA GGT TGA TAA CGC TCC AGA AGA-3'		
RpoB-F	5'-CGW ATG AAC ATY GGB CAG GT-3'	406	4
RpoB-R	5'-TCC ATY TCR CCR AAR CGC TG-3'		

\*1: ×1 (95 °C, 600 s), ×30 (95 °C, 30 s, 58 °C, 60 s, 72 °C, 60 s), and ×1 (72 °C, 420 s). 2: ×1 (94 °C, 180 s), ×30 (94 °C, 30 s, 50 °C, 40 s, 72 °C, 60 s), and ×1 (72 °C, 300 s). 3: ×1 (94 °C, 180 s), ×30 (94 °C, 45 s, 57 °C, 40 s, 72 °C, 60 s), and ×1 (72 °C, 420 s). 4: ×1 (95 °C, 600 s), ×35 (94 °C, 30 s, 50 °C, 30 s, 72 °C, 120 s), and ×1 (72 °C, 600 s)

**Table 2** Biochemical properties of *A. pinnipediorum* 014418 of the present study and type strain *A. pinnipediorum* DSM 28752<sup>T</sup> using VITEK2-compact system


Test*	<i>A. pinnipediorum</i> 014418	<i>A. pinnipediorum</i> DSM 28752 <sup>T</sup>
D-GALACTOSE	+	–
D-CELLOBIOSE	–	–
SACCHAROSE/SUCROSE	–	–
BETA-GALACTOPYRANOSIDASE Indoxyl	+	+
MALTOTRIOSE	–	–
PHOSPHATASE	–	–
Leucine-ARYLAMIDASE	+	+
Tyrosine-ARYLAMIDASE	+	+
ARGININ-GP	–	–
ALPHA-ARABINOSIDASE	–	–
AESCULIN-Hydrolyse	–	–
L-ARABINOSE	–	–
ELLMAN	–	–
Ala-Phe-Pro-ARYLAMIDASE	+	+
N-ACETYL-D-GLUCOSAMIN	+	+
5-Brom-4-Chlor-3-Indoxyl-alpha-Galactoside	–	–
BETA-D-FUCOSIDASE	–	+
d-Ribose 2	–	–
Phenylalanin-ARYLAMIDASE	+	+
D-GLUCOSE	+	+
5-Brom-4-Chlor-3-Indoxyl-beta-Glucoside	–	–
BETA-MANNOSIDASE	–	–
5-Brom-4-Chlor-3-Indoxyl-beta-N-Acetyl-Glucosamide	+	+
Phenylphosphonate	–	–
L-Prolin-ARYLAMIDASE	+	+
D-MANNOSE	–	–
UREASE	–	–
5-Brom-4-Chlor-3-Indoxyl-alpha-Mannoside	–	–
ALPHA-L-ARABINFURANOSIDE	–	–
L-Pyrrolidonyl-ARYLAMIDASE	+	+
D-MALTOSE	–	–
5-Brom-4-Chlor-3-Indoxyl-beta-Glucuronide	–	–
PYRUVAT	–	–
ALPHA-L-FUCOSIDASE	–	–
D-XYLOSE	–	–
Gram-stain	+	+
Morph (Morphology)	–	–
Aero (Aerotolerance)	+	+

\*Tests of VITEK2-compact (bioMérieux, Nürtingen, Germany). The reactions are shown as follows: + = positive reaction; – = negative reaction

014418 could additionally be identified to the species level by MALDI-TOF MS analysis with the current Bruker database (MBT 8468 MSP library). *A. pinnipediorum* 014418 displayed a close relation to the type strain *A. pinnipediorum* DSM 28752<sup>T</sup> and could be clearly separated from other species of genus *Arcanobacterium*. A typical dendrogram of the MALDI-TOF mass spectrum analysis of *A. pinnipediorum* 014418 and various other species of genus *Arcanobacterium*

is shown in Fig. 1. MALDI-TOF MS has already been shown to be a rapid and reliable technique for identification of bacteria of genus *Arcanobacterium* and *Trueperella* (Hijazin et al. 2012a; Wickhorst et al. 2017), also including *A. phocae* (Alssahen et al. 2018), *A. phocisimile* (Sammra et al. 2014a), and *A. pinnipediorum* (Sammra et al. 2018) of seal origin. MALDI TOF mass-spectra of *A. pinnipediorum* 014418 and other species of genus *Arcanobacterium* are available by

**Fig. 1** Dendrogram of MALDI-TOF MS main spectra of *A. pinnipediorum* 014418 investigated in the present study in comparison with type strain *A. pinnipediorum* DSM 28752<sup>T</sup> and other species of genus *Arcanobacterium*. The MALDI-TOF MS analysis was performed using MALDI Biotyper Version (4.0)



exchange via the MALDI-TOF user platform (<https://www.maldi-up.ua-bw.de>; Rau et al. 2016).

FT-IR spectroscopy has already been used as a tool for identification of a large number of clinically relevant pathogens (Kuhm et al. 2009; Samuels et al. 2009; Contzen et al. 2011; Grunert et al. 2013), also including *T. pyogenes* isolated from bovine mastitis (Nagib et al. 2014) and for characterization of *A. pinnipediorum* of seal origin (Sammra et al. 2018). The infrared spectra of *A. pinnipediorum* 014418 of the present study was analyzed by the method described by Nagib et al. (2014). Comparable to the MALDI-TOF MS analysis *A. pinnipediorum* 014418 of the present study yielded a close relation to *A. pinnipediorum* DSM 28752<sup>T</sup> and to next closely related members *A. phocae* DSM 10002<sup>T</sup>, *A. phocisimile* DSM 26142<sup>T</sup>, and *A. haemolyticum* DSM 20595<sup>T</sup> (Fig. 2).


The previously described loop mediated isothermal amplification (LAMP) assay for detection of phocaelysin encoding gene *phl* could be used successfully to identify *A. phocae* of various origins (Alssahen et al. 2018). However, *A. pinnipediorum* 014418 and comparatively investigated *A. pinnipediorum* DSM 28752<sup>T</sup> and *A. phocisimile*


DSM 26142<sup>T</sup> yielded a negative *phl*-reaction indicating the high specificity of this assay for detection of *A. phocae* but not for other species of genus *Arcanobacterium* of seal origin (data not shown).

Strain *A. pinnipediorum* 014418 of the present study was identified genotypically by sequencing the 16S rRNA gene with a sequence similarity of 99.9% to 16S rRNA gene of type strain *A. pinnipediorum* DSM 28752<sup>T</sup> (KJ596349) and a 16S rRNA gene sequence similarity  $\leq 98.7\%$  to other species of genus *Arcanobacterium* (Fig. 3).

According to previous studies *A. phocae*, *A. phocisimile* and *A. pinnipediorum* of seal origin could be further characterized genotypically by sequencing the genomic targets *gap*, *tuf*, and *rpoB* (Sammra et al. 2014a, 2018). Comparable to these studies, *A. pinnipediorum* 014418 showed sequence similarities of the genes *gap*, *tuf*, and *rpoB* of 97.1%, 98.7%, and 93.6%, respectively, to type strain *A. pinnipediorum* DSM 28752<sup>T</sup> and sequence similarities of these genes of  $\leq 91.2\%$ ,  $\leq 94.0\%$ , and  $\leq 86.5\%$ , respectively, to other species of genus *Arcanobacterium*. A dendrogram analysis of these target genes is shown in Fig. 4a, b, c.

**Fig. 2** Cluster analysis of infrared spectra of *A. pinnipediorum* 014418 investigated in the present study in comparison with type strain *A. pinnipediorum* DSM 28752<sup>T</sup> and other species of genus *Arcanobacterium*. This analysis was performed by using the second derivatives of the spectra ( $n=2$  for each isolate) in the spectral range of 500 to 1400  $\text{cm}^{-1}$ . Ward's algorithm was applied in OPUS (version 7.2, Bruker optic, Ettlingen)






**Fig. 3** Dendrogram analysis of 16S rRNA gene of *A. pinnipediorum* 014418 investigated in the present study in comparison with the type strain *A. pinnipediorum* DSM 28752<sup>T</sup> and type strains of other species of genus *Arcanobacterium* obtained from NCBI GenBank using

the Clustal W method of DNASTAR/Lasergene MegAlign program (version 8.0.2). \*Accession numbers are given in brackets. <sup>T</sup> indicates type strains


According to the results of the present study, *A. pinnipediorum* 014418 isolated from mandibular discharge of a grey seal pup in England (UK) could successfully be

characterized biochemically, by MALDI-TOF MS, by FT-IR spectroscopy, and by sequencing the presented genomic targets. The usefulness of the determination of specific spectra


**Fig. 4** Dendrogram analyses of the genes *gap a*, *tuf b*, and *rpoB c* of *A. pinnipediorum* 014418 investigated in the present study in comparison with the type strain *A. pinnipediorum* DSM 28752<sup>T</sup> and type strains of other species of genus *Arcanobacterium* obtained from NCBI GenBank using the Clustal W method of DNASTAR/Lasergene MegAlign program (version 8.0.2). \*Accession numbers are given in brackets



**(a) gap**



**(b) tuf**



**(c) rpoB**



by MALDI-TOF MS and FT-IR analyses and the various genotypic targets for identification of this species have to be further investigated with additional *A. pinnipediorum* strains isolated in other contexts. The biochemical differences observed between *A. pinnipediorum* 014418 and the type strain *A. pinnipediorum* DSM 28752<sup>T</sup> might reflect an intra-specific variability caused by evolutionary processes. As already mentioned above, further phenotypical studies and whole genome sequence analyses of *A. pinnipediorum* of various origins will show the unique taxonomic position and the importance of this species in animal infections. However, strain *A. pinnipediorum* 014418 of the present study represented the first isolation of this species in the UK and the first isolation from a grey seal. *A. pinnipediorum* 014418 was isolated from a mixed culture with several other bacteria, including *Enterococcus faecalis*, *Streptococcus lutetiensis*, and a Gram negative cocco-bacillus that could not be identified further, indicating that the pathogenic importance of this species remains to be elucidated. It is perhaps of note, however, that *A. phocae* has often been isolated in mixed infections, including, as is the case in this report, *Streptococcus* spp., from seals in Scotland (G. Foster, unpublished findings). Furthermore, *A. phocae* has been recovered from local lesions including an infected jaw from a mouth lesion of a grey seal (Ramos et al. 1997), providing further similarities with the isolation of *A. pinnipediorum* 014418 from an infected jaw.

**Author contribution** M.A., C.L., G.F., A.A., T.E., A.H., E.P-B, and J.R. contributed to the design of the study and collected and analyzed the data. M.A. performed the initial examination of the isolates. M.A., C.L., A.A., and G.F. drafted the manuscript. All authors read and approved the final manuscript.

**Funding** Open Access funding enabled and organized by Projekt DEAL.

## Declarations

**Conflict of interest** The authors declare no competing interests.

**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

## References

- Abdulmawjood A, Wickhorst J, Hashim O, Sammra O, Hassan AA, Alssahen M, Lämmler C, Prenger-Berninghoff E, Klein G (2016) Application of a loop-mediated isothermal amplification (LAMP) assay for molecular identification of *Trueperella pyogenes* isolated from various origins. *Mol Cell Probes* 30:205–210
- Alssahen M, Sammra O, Wickhorst J, Hassan AA, Lämmler C, Raunio-Saarnisto M, Prenger-Berninghoff E, Timke M, Becker A, Abdulmawjood A (2018) Identification of *Arcanobacterium phocae* isolated from fur animals by phenotypic properties, by MALDI-TOF MS analysis and by detection of phocaelysin encoding gene *phl* as probable novel target. *Vet Microbiol* 216:45–51
- Collins MD, Jones D, Schofield GM (1982) Reclassification of *Corynebacterium haemolyticum* (MacLean, Liebow & Rosenberg) in the genus *Arcanobacterium* gen. nov. as *Arcanobacterium haemolyticum* nom.rev., comb.nov. *J Gen Microbiol* 28:1279–1281
- Contzen M, Sting R, Blazey B, Rau J (2011) *Corynebacterium ulcerans* from diseased wild boars. *Zoonoses Public Health* 58:479–488
- Diop K, Morand A, Dubus JC, Fournier EP, Raoult D, Fenollar F (2017) *Arcanobacterium urinimassiliense* sp. nov., a new bacterium isolated from the urogenital tract. *New Microbe New Infect* 18:15–17
- Fall NS, Lo CI, Fournier PE, Sokhna C, Raoult D, Fenollar F, Lagier JC (2019) *Arcanobacterium ihumii* sp. nov., *Varibaculum vaginae* sp.nov. and *Tessaracoccus timonensis* sp. nov., isolated from vaginal swabs from healthy Senegalese women. *New Microbe New Infect* 31:100585
- Grunert T, Wenning M, Barbagelata MS, Fricker M, Sordelli DO (2013) Rapid and reliable identification of *Staphylococcus aureus* capsular serotypes by means of artificial neural network-assisted Fourier transform infrared spectroscopy. *J Clin Microbiol* 51:2261–2266
- Hassan AA, Ülbegi-Mohyla H, Kanbar T, Alber J, Lämmler C, Abdulmawjood A, Zschöck M, Weiss R (2009) Phenotypic and genotypic characterization of *Arcanobacterium haemolyticum* isolates from infections of horses. *J Clin Microbiol* 47:124–128
- Hijazin M, Hassan AA, Alber J, Lämmler C, Timke M, Kostorzewa M, Prenger-Berninghoff E, Zschöck M (2012a) Evaluation of matrix-assisted laser desorption/ionization-time-of-flight mass spectrometry (MALDI-TOF MS) for species identification of bacteria of genera *Arcanobacterium* and *Trueperella*. *Vet Microbiol* 157:243–245
- Hijazin M, Prenger-Berninghoff E, Sammra O, Alber J, Lämmler C, Kämpfer P, Glaeser SP, Busse HJ, Hassan AA, Abdulmawjood A, Zschöck M et al (2012b) *Arcanobacterium canis* sp. nov., isolated from otitis externa of a dog, and emended description of the genus *Arcanobacterium* Collins, 1983 emend. Yassin et al., 2011. *Int J Syst Evol Microbiol* 62:2201–2205
- Hijazin M, Sammra O, Ülbegi-Mohyla H, Nagib S, Alber J, Lämmler C, Kämpfer P, Glaeser SP, Busse HJ, Kassmannhuber J, Prenger-Berninghoff E, Weiss R, Siebert U, Hassan AA, Abdulmawjood A, Zschöck M (2013) *Arcanobacterium phocisimile* sp. nov., isolated from harbour seals. *Int J Syst Evol Microbiol* 63:2019–2024
- Kuhm AE, Suter D, Felleisen R, Rau J (2009) Identification of *Yersinia enterocolitica* at the species and subspecies levels by Fourier transform infrared spectroscopy. *Appl Environ Microbiol* 75:5809–5813
- Nagib S, Rau J, Sammra O, Lämmler C, Schlez K, Zschöck M, Prenger-Berninghoff E, Klein G, Abdulmawjood A (2014) Identification of *Trueperella pyogenes* isolated from bovine mastitis by Fourier transform infrared spectroscopy. *PLoS One* 9:e104654

- Ramos C, Foster G, Collins MD (1997) Phylogenetic analysis of the genus *Actinomyces* based on 16S rRNA gene sequences: description of *Arcanobacterium phocae* sp. nov., *Arcanobacter bernardiae* comb. nov. and *Arcanobacterium pyogenes* comb. nov. J Clin Microbiol 47:46–53
- Rau J, Eisenberg T, Männig A, Wind C, Lasch R, Sting R (2016) MALDI-UP – An internet platform for the exchange of mass spectra — user guide for <http://maldi-up.ua-bw.de/>. Aspects of food control and animal health 01/2016. <http://ejournal.cvuas.de/issue201601.asp>
- Sammra O, Balbutskaya A, Zhang S, Hijazin M, Nagib S, Lämmle C, Abdulmawjood A, Prenger-Berninghoff E, Kostrzewa M, Timke M (2013) Further characteristics of *Arcanobacterium canis*, a novel species of genus *Arcanobacterium*. Vet Microbiol 167:619–622
- Sammra O, Balbutskaya A, Hijazin M, Nagib S, Alber J, Lämmle C, Abdulmawjood A, Prenger-Berninghoff E, Timke M, Kostrzewa M, Siebert U (2014a) Further study on *Arcanobacterium phocisimile* a novel species of genus *Arcanobacterium*. J Vet Med ID 923592
- Sammra O, Balbutskaya A, Nagib S, Alber J, Lämmle C, Abdulmawjood A, Timke M, Kostrzewa M, Prenger-Berninghoff E (2014b) Properties of an *Arcanobacterium haemolyticum* strain isolated from a donkey. Berl Münch Tierärztl Wochenschr 127:10–14
- Sammra O, Balbutskaya A, Ülbegi-Mohyla H, Nagib S, Lämmle C, Kämpfer P, Glaeser SP, Golke J, Busse HJ, Prenger-Berninghoff E, Siebert U, Abdulmawjood A, Klein G (2015) *Arcanobacterium pinnipediorum* sp. nov., isolated from a harbour seal. Int J Syst Evol Microbiol 65:4539–4543
- Sammra O, Foster G, Hassan AA, Alssahen M, Lämmle C, Glaeser SP, Kämpfer P, Busse H-J, Borowiak M, Malorny B, Ritchie CM, Prenger-Berninghoff E, Abdulmawjood A (2020) *Arcanobacterium bovis* sp. nov., isolated from the milk of a cow with mastitis. Int J Syst Evol Microbiol 70:4105–4110. <https://doi.org/10.1099/ijsem.0.004230>
- Sammra O, Rau J, Wickhorst JP, Alssahen M, Hassan AA, Lämmle C, Kämpfer P, Glaeser SP, Busse HJ, Kleinhagauer T, Knauf-Witzens T, Prenger-Berninghoff E, Abdulmawjood A, Klein G (2017) *Arcanobacterium wilhelmae* sp. nov., isolated from the genital tract of a rhinoceros (*Rhinoceros unicornis*). Int J Syst Evol Microbiol 67:2093–2097
- Sammra O, Rau J, Wickhorst JP, Alssahen M, Hassan AA, Lämmle C, Prenger-Berninghoff E, Abdulmawjood A (2018) Further characteristic of *Arcanobacterium pinnipediorum* DSM 28752<sup>T</sup> and *Arcanobacterium wilhelmae* DSM 102162<sup>T</sup>, two novel species of genus *Arcanobacterium*. Folia Microbiol 63:695–700
- Samuels AC, Snyder AP, Emge DK, Amant D, Minter J (2009) Classification of select category A and B bacteria by Fourier transform infrared spectroscopy. Appl Spectrosc 63:14–24
- Ülbegi-Mohyla H, Hijazin M, Alber J, Lämmle C, Hassan AA, Abdulmawjood A, Prenger-Berninghoff E, Weiss R, Zschöck M (2010) Identification of *Arcanobacterium pyogenes* isolated by post mortem examinations of a bearded dragon and a gecko by phenotypic and genotypic properties. J Vet Sci 11:265–267
- Wickhorst JP, Hassan AA, Sheet OH, Eisenberg T, Sammra O, Alssahen M, Lämmle C, Prenger-Berninghoff E, Zschöck M, Timke M, Abdulmawjood A (2017) *Trueperella pyogenes* isolated from a brain abscess of an adult roebuck (*Capreolus capreolus*). Folia Microbiol 63:17–22
- Wickhorst JP, Hassan AA, Sammra O, Alssahen M, Lämmle C, Prenger-Berninghoff E, Naggert M, Timke M, Rau J, Abdulmawjood A (2019) First report on the isolation of *Trueperella abortus* from companion animals. Res Vet Sci 125:465–467
- Yassin AF, Hupfer H, Siering C, Schumann P (2011) Comparative chemotaxonomic and phylogenetic studies on the genus *Arcanobacterium* Collins et al., 1982 emend. Lehnert et al., 2006: proposal for *Trueperella* gen. nov. and emended description of the genus *Arcanobacterium*. Int J Syst Evol Microbiol 61:1265–1274

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.